arXiv:cond-mat/9706149v2 21 Aug 1997

Baryon Asymmetry of Universe: View from Superfluid ^{3}He

G.E. Volovik *

Helsinki University of Technology, Low Temperature Laboratory, P.O.Box 2200,
FIN-02015 HUT, Finland
**permanent address: Landau Institute for Theoretical Physics, Moscow, Russia*

The origin of the excess of matter over antimatter in our Universe remains one of the fundamental problems. Dynamical baryogenesis in the process of the broken symmetry electroweak transition in the expanding Universe is the widely discussed model where the baryonic asymmetry is induced by the quantum chiral anomaly. We discuss the modelling of this phenomenon in superfluid ^{3}*He and superconductors where the chiral anomaly is realized in the presence of quantized vortex, which introduces nodes into the energy spectrum of the fermionic quasiparticles. The spectral flow of fermions through the nodes during the vortex motion leads to the creation of fermionic charge from the superfluid vacuum and to transfer of the superfluid linear momentum into the heat bath, thus producing an extra force on the vortex, which in some cases compensates the Magnus force. This spectral-flow force was calculated 20 years ago by Kopnin and Kravtsov for s-wave superconductors, but only recently was it measured in a broad temperature range in Manchester experiments on rotating superfluid* ^{3}*He. The "momentogenesis" observed in* ^{3}*He is analogous to the dynamical production of baryons by cosmic strings. Some other possible scenaria of baryogenesis related to superfluid* ^{3}*He are discussed.*

PACS numbers: 67.57.-z, 11.27.+d, 98.80.Cq, 11.30.Fs

1. INTRODUCTION

The superfluid phases of ^{3}He and the electroweak vacuum of the standard Weinberg-Salam model share many common properties[1] (see Fig.1). The most important similarity is that the fermionic quasiparticles in superfluid ^{3}He interact with the order parameter in a way that is similar to the interaction of fermions with the electroweak gauge and Higgs fields. Within this analogy the baryon number violation appears to be analogous to the violation of fermion momentum conservation within an A-phase vortex. The phenomenon which unites the barygenesis in the Weinberg-Salam model (electroweak baryogenesis) and the momentogenesis in the pair-correlated fermionic systems is known generically as the "chiral anomaly". There are certain quantities, of which baryon number is an example, which are classically conserved but can be violated by quantum mechanical effects. The process leading to particle creation is called "spectral flow", and can be pictured as a process in which fermions flow under an external perturbation from negative energy levels towards positive energy levels. Some fermions therefore cross zero energy and move from the Dirac sea (vacuum) into the observable positive energy world (matter), introducing an extra charge of the matter.

The analogous process in superfluids/superconductors is the transfer of the linear (or angular) momentum from the condensate (vacuum) to the heat bath of the normal excitations (matter). Thus the momentum of the "matter" is not conserved, while the conservation of the total momentum of vacuum+matter is not violated. The striking feature of this analogy is that the rate of momentum transfer due to the spectral flow is described by the same equation of the chiral anomaly which was introduced by Adler[2] and Bell and Jackiw[3] in quantum field theory. This is vusualized on the basis of the vortex texture in ^{3}He-A. However, such anomalous violation of fermionic charge is a general phenomenon in condensed matter and occurs also for vortices in ^{3}He-B and Abrikosov vortices in superconductors.

1.1. Chiral Anomaly.

It is known that the charge particle with definite chirality, say, right-handed electron, has a peculiar property in the presence of magnetic field. The Landau quantization of the motion of this particle in magnetic field $\mathbf{B} \parallel \hat{z}$ leads to an anomalous gapless branch $E(p_z) = cp_z$, which crosses zero energy at $p_z = 0$ (Fig.2). This branch is asymmetric with respect to the spatial inversion $p_z \to -p_z$. As a result the vacuum state is asymmetric since only the states with negative p_z are occupied. The spectral flow of the electrons from the vacuum state to the positive energy world leads to the nucleation of the matter from the vacuum. This occurs if the electric field is applied along z. According to the second Newton law the linear momentum of the particle increases linearly with time, $\partial_t p_z = e_R E_z$, where e_R is the electric charge of right electron. This means that under applied field the particles from negative energy levels of Dirac vacuum are pushed to positive energy levels of matter with the rate determined by the electric field projection E_z on the magnetic field and by the density of state $|e_R\mathbf{B}|/(2\pi)^2$ on the lowest Landau level:

$$\partial_t n = \frac{1}{4\pi^2} e_R^2 \mathbf{E}\cdot\mathbf{B} \tag{2.1.1a}$$

The left-handed electrons have the anomalous branch with the opposite slope $E(p_z) = -cp_z$, as a result the production of the particles from the vacuum has the opposite sign:

$$\partial_t n = -\frac{1}{4\pi^2} e_L^2 \mathbf{E}\cdot\mathbf{B} \tag{2.1.1b}$$

where e_L is the charge of the left electron. If the parity is conserved, then $e_L = e_R$, and the net production of the particles from the vacuum is zero. In electroweak interactions and in ^{3}He-A the parity is broken, and this leads to the chiral anomaly: uncompensated anomalous production of the charge from the vacuum.

1.2. Baryon Production by Chiral Anomaly.

In the Weinberg-Salam model there are two types of magnetic and electric fields: hypercharge fields $\mathbf{B}_Y$ and $\mathbf{E}_Y$ and weak fields $\mathbf{B}_W$ and $\mathbf{E}_W$. The hypercharge Y, and weak charge W of left and right quarks are essentially different:

$$Y_{dR} = -\frac{1}{3},\ Y_{uR} = \frac{2}{3},\ W_{dL} = -W_{uL} = \frac{1}{2},\ W_{dR} = W_{uR} = 0,\ Y_{dL} = Y_{uL} = \frac{1}{6} \tag{2.2.1}$$

Here the subscripts uL and uR mean the left and right u-quarks; the same is for the d-quarks. The total number of the u- and d- quarks produced by the electric and magnetic fields per unit time per unit volume is thus

$$\frac{3}{4\pi^2}[\mathbf{B}_Y\cdot\mathbf{E}_Y(Y_{dR}^2+Y_{uR}^2-Y_{dL}^2-Y_{uL}^2)+\mathbf{B}_W\cdot\mathbf{E}_W(W_{dR}^2+W_{uR}^2-W_{dL}^2-W_{uL}^2)]\ , \tag{2.2.2}$$

where the factor 3 appears due to 3 colours of quarks. Substituting all the charges from Eq.(2.2.1) and taking into account that there are $N_F = 3$ families of quarks and that all the quarks have the same baryon number $B = 1/3$ one obtains the following rate of the baryoproduction

$$\partial_t B = \frac{N_F}{8\pi^2}\left(-\mathbf{B}_W\cdot\mathbf{E}_W + \mathbf{B}_Y\cdot\mathbf{E}_Y\right)\ . \tag{2.2.3}$$

The nonconservation of B is thus the result of the phenomenon of the chiral (axial) anomaly predicted by Adler[2] and Bell and Jackiw.[3] The fields $\mathbf{B}_Y$, $\mathbf{E}_Y$, $\mathbf{B}_W$ and $\mathbf{E}_W$ can be generated in the core of topological defects, such as monopoles, domain walls, sphalerons and electroweak cosmic strings evolving in the expanding Universe.[4,5,6,7,8,9]

1.3. Momentogenesis in 3He-A.

The anomaly mediated transfer of the linear momentum can be visualized through the example of the continuous distribution of the vector $\hat{\mathbf{l}}$ in ^{3}He-A, which shows the direction in the momentum space, where the spectrum of quasiparticles has a node. In the vicinity of the point gap nodes the quasiparticles are chiral: like the neutrino they are either left-handed or right-handed.[10] The interaction of the quasiparticles with the $\hat{\mathbf{l}}$ texture is characterized by the "vector potential" $\mathbf{A} = k_F\hat{\mathbf{l}}$, where $p_F = \hbar k_F$ is the magnitude of the momentum at the position of the node. For such gapless fermions the anomaly in Eq.(2.1.1) takes place producing a "chiral charge" of quasiparticles with the rate $(1/2\pi^2)\,\mathbf{E}\cdot\mathbf{B} = (k_F^2/2\pi^2)\,\partial_t\hat{\mathbf{l}}\cdot(\vec{\nabla}\times\hat{\mathbf{l}})$ where the *electric* and *magnetic* fields are $\mathbf{E} = \partial_t\mathbf{A}$, $\mathbf{B} = \vec{\nabla}\times\mathbf{A}$. The linear momentum produced by the chiral anomaly is

$$\partial_t\mathbf{P} = \frac{1}{4\pi^2}(\mathbf{B}\cdot\mathbf{E})\,[\mathbf{P}_R - \mathbf{P}_L] \quad . \qquad (2.3.1)$$

where $\mathbf{P}_R$ and $\mathbf{P}_L$ – momenta of right and left quasiparticle. The Left-Right asymmetry, $\mathbf{P}_R = -\mathbf{P}_L = p_F\hat{\mathbf{l}}$, gives the net momentoproduction

$$\partial_t\mathbf{P} = \frac{1}{2\pi^2}\int d^3r\; p_F\hat{\mathbf{l}}\;(\mathbf{B}\cdot\mathbf{E}) = \hbar\frac{k_F^3}{2\pi^2}\int d^3r\;\hat{\mathbf{l}}\;(\partial_t\hat{\mathbf{l}}\cdot(\vec{\nabla}\times\hat{\mathbf{l}}\;)) \quad . \qquad (2.3.2)$$

Since the total linear momentum is conserved in condensed matter, the Eq.(2.3.2) means that, in the presence of a time-dependent texture, the momentum is transferred from the superfluid *vacuum* to the *matter* (the system of the excitations).

1.4. Momentogenesis by a Continuous Vortex

The simplest time-dependent texture, which is experimentally realized in ^{3}He-A, is the moving continuous vortex. The latter corresponds to topologically unstable continuous strings in electroweak theory.[1] The simplest type of continuous vortex has the following distribution of $\hat{\mathbf{l}}$-field and superfluid velocity $\mathbf{v}_s$ in cylindrical coordinate system:

$$\hat{\mathbf{l}}(r,\phi) = \hat{\mathbf{z}}\cos\eta(r) + \hat{\mathbf{r}}\sin\eta(r)\ ,\ \mathbf{v}_s(r,\phi) = -\frac{\hbar}{2mr}[1+\cos\eta(r)]\hat{\phi} \quad , \quad (2.4.1)$$

where $\eta(r)$ changes from $\eta(0) = 0$ to $\eta(\infty) = \pi$ in the so called soft core of the vortex. The stationary vortex generates the "magnetic" field $\mathbf{B} = k_F\vec{\nabla}\times\hat{\mathbf{l}}$. When the vortex moves with a constant velocity $\mathbf{v}_L$ (with respect to the heat bath) it also generates the "electric" field, since $\mathbf{A}$ depends on $\mathbf{r} - \mathbf{v}_L t$:

$$\mathbf{E} = \partial_t\mathbf{A} = -k_F(\mathbf{v}_L\cdot\vec{\nabla})\hat{\mathbf{l}} \quad . \qquad (2.4.2)$$

Integration of the anomalous momentum transfer in Eq.(2.3.2) over the cross-section of the soft core of the moving vortex gives the rate of the momentum transfer between the condensate and the heat bath, mediated by the moving vortex:[11]

$$\partial_t \mathbf{P} = -\pi\hbar N_v C_0 \hat{\mathbf{z}} \times \mathbf{v}_L \quad . \tag{2.4.3}$$

Here N_v is the winding number of the vortex ($N_v = -2$ in the case of Eq.(2.4.1)) and $C_0 = k_F^3/3\pi^2$.

This corresponds to an extra nondissipative force $\mathbf{F}_{sf} = \partial_t \mathbf{P}$ acting on the moving continuous vortex, which is called the spectral-flow force. This result for the continuous vortex, derived from the axial anomaly equation (2.1.1), was confirmed by the microscopic theory, which took into account the discreteness of the quasiparticle spectrum in the soft core.[12]

1.5. Spectral-flow force vs Magnus force.

The spectral-flow force is the transverse nondissipative force which acts on the vortex, when it moves with respect to the matter (heat bath or normal component):

$$\mathbf{F}_{sf} = -\pi\hbar N_v C_0 \hat{\mathbf{z}} \times [\mathbf{v}_L - \mathbf{v}_n] \quad , \tag{2.5.1}$$

where $\mathbf{v}_n$ is the velocity of the normal component of the liquid. In addition to this there is also an old classical Magnus force, which acts on the vortex moving with respect to superflow:

$$\mathbf{F}_M = \pi\hbar N_v \frac{\rho}{m} \hat{\mathbf{z}} \times [\mathbf{v}_L - \mathbf{v}_s] \quad , \tag{2.5.2}$$

where $\mathbf{v}_s$ is the superfluid velocity of the coherent condensate (vacuum) and ρ/m is the particle density. The balance of these two forces acting on the vortex (together with the Iordanskii force, which we do not discuss here, see Ref.,[13] and the longitudinal friction force, see below in Sec. 3.2) determines the motion of the vortex and thus the mutual force between the vacuum and matter mediated by the moving vortex.

It is important, that if $\mathbf{v}_n = \mathbf{v}_s$ the spectral flow almost completely compensates the Magnus force. This is because the particle density ρ/m and the spectral-flow parameter $C_0 = k_F^3/3\pi^2$ are very close to each other. In fact in the normal Fermi liquid above the superfluid transition temperature T_c these quantities coincide: $\rho(T > T_c) = mk_F^3/3\pi^2$. In the superfluid phase the disbalance between ρ/m and C_0 occurs due to the tiny asymmetry between particles and holes: $\rho(T < T_c) - mk_F^3/3\pi^2 \sim \rho(\Delta(T)/E_F)^2 \ll \rho$, where $\Delta(T)$ is the gap amplitude and E_F the Fermi-energy. Such cancellation of the Magnus force was confirmed in experiments on the vortex dynamics in ^{3}He-A.[14,15]

2. MOMENTOGENESIS BY SINGULAR VORTICES

The momentum is transferred from the vacuum to the heat bath by the vortex motion, the chiral anomaly being the main mechanism which determines the rate of momentum transfer. However, in the case of vortices with a singular core (in superconductors and in ^{3}He-B) the equation (2.1.1) for the axial anomaly is no more valid: the anomalous force should be calculated directly from the consideration of the spectral flow in the vortex core. This is similar to the calculation of the baryonic charge on Z-strings, where instead of the anomaly equations one must operate with the exact spectrum of fermions in the string core.[9]

2.1. Spectral Flow in Singular Vortex.

The spectrum of single-fermion excitations has anomalous (chiral) branch $E(p_z, L_z)$, which depends on the momentum projection p_z on the vortex axis and the discrete angular momentum L_z in the following way:[12]

$$E(p_z, L_z) = -\omega_0(p_z)\ (L_z - L_0(p_z)) \qquad (3.1.1)$$

For most symmetric vortices, where $L_0(p_z) = 0$, the spectrum was first obtained in.[16] The interlevel distance $\hbar\omega_0(p_z)$, the so called minigap, is small compared to the gap amplitude $\Delta(T)$ of the fermions outside the core: $\hbar\omega_0 \sim \Delta^2(T)/E_F \ll \Delta(T)$, where again E_F is the Fermi-energy. As a function of L_z, the anomalous branch crosses zero energy (Fig.3). According to the index theorem, the number of such anomalous branches equals the winding number N_v of the vortex[17]). This results in a momentum flow from the the Dirac sea to the heat bath, if the minigap $\hbar\omega_0$ between the core states can be neglected. We consider just this case, which occurs when ω_0 is less than the inverse relaxation time, $1/\tau$.

If the velocity v_L is along x, the excitation angular momentum L_z in the moving core changes with the rate $\dot L_z = \dot x p_y = v_L p_y$. Since the neighbouring levels differ by $\Delta L_z = \hbar$ and the number of anomalous branches equals the winding number N_v, the levels cross zero at a rate $-N_v \dot L_z/\hbar$. Thus the flow of the momentum p_y from the negative to the positive energy states occurs with the rate $\dot p_y = -N_v p_y \dot L_z/\hbar = -N_v v_L p_y^2/\hbar$. Integrating this momentum transfer over the Fermi surface and taking into account that the in-plane average $< p_y^2 >= \frac{1}{2}(p_F^2 - p_z^2)$, one again obtains the Eq.(2.4.3)

$$\partial_t \mathbf{P} = \hat{\mathbf{y}}\partial_t P_y = -\frac{1}{2\hbar}\hat{\mathbf{y}} v_L N_v \int_{-p_F}^{p_F} \frac{dp_z}{2\pi\hbar}(p_F^2 - p_z^2) = -N_v \pi\hbar C_0 \hat{\mathbf{z}}\times\mathbf{v}_L \ . \qquad (3.1.2)$$

Thus the result of the direct calculation of the spectral flow in the singular core coincides with the result obtained for the continuous ^{3}He-A vortex from the axial anomaly equation. This is not a coincidence, but the consequence of the underlying topology in the 6-dimensional real+momentum space. The singular vortex in any Fermi superfluid/superconductor can be transformed into a continuous one by extending the core. The order parameter field in the extended core is similar to that of ^{3}He-A: the spectrum of quasiparticles contains the point nodes.[20] In the vicinity of the nodes the quasiparticles are chiral leading to the Adler-Bell-Jackiw anomaly and thus to the spectral-flow force.

2.2. Nondissipative and Friction Forces.

The anomalous nondissipative force acting on the vortex has maximal value if the spectral flow is not suppressed by the minigap, i.e. at $\omega_0\tau \ll 1$. In the general case considered in[18,19] and experimentally measured in superfluid ^{3}He-B,[14,15] the spectral flow force is reduced by the factor $1/(1+\omega_0^2\tau^2)$ and also the dissipative friction component of the force appears:

$$\mathbf{F}_{sf} = N_v \pi\hbar C_0 \left[\frac{1}{1+\omega_0^2\tau^2}\hat{\mathbf{z}} \times (\mathbf{v}_n - \mathbf{v}_L) + \frac{\omega_0\tau}{1+\omega_0^2\tau^2}(\mathbf{v}_n - \mathbf{v}_L)\right] \quad , \quad (3.2.1)$$

where $\mathbf{v}_n$ is the velocity of the heat bath (or velocity of the normal component of the superfluid/superconductor). This expression was found more than 20 years ago by Kopnin and Kravtsov in the Gorkov formalism developed for s-wave superconductivity.[21] In the d-wave superconductor the situation is more interesting due to the gapless quasiparticles outside the vortex cores.[22] In this case the minigap ω_0 depends on the orientation and has nodes. This leads to the finite dissipation even in the superclean limit $< \omega_0 > \tau = \infty$, which occurs due to the Landau damping. In the moderately clean regime the scaling law for the flux flow at low T and low field is obtained. The gapless regime can also occur for ^{3}He-B vortices with the spontaneously broken continuous symmetry of the core, this leads to effects corresponding to the Hawking radiation at the event horizon in black holes.[23,24]

3. TOPOLOGY OF MOMENTOGENESIS

3.1. Topological and Fermionic Charges

The vortex is the object which serves as a mediator between the condensate (vacuum) and heat bath (matter) and which allows them to exchange linear momentum, while the total momentum is conserved:

$$\mathbf{P} = \mathbf{P}_{vac} + \mathbf{P}_{matter} = \int d^3r \; \rho \mathbf{v}_s + \sum_{\mathbf{p}} f(\mathbf{p})\mathbf{p} \quad . \quad (4.1.1)$$

Here $\mathbf{v}_s$ is the superfluid velocity which in terms of the phase Φ of the order parameter is $\mathbf{v}_s = \frac{\kappa}{2\pi}\vec{\nabla}\Phi$; κ is the circulation quantum, $\kappa = \pi\hbar/m_3$ for the superfluid ^{3}He, with m_3 being is the mass of ^{3}He atom); ρ is the mass density and $f(\mathbf{p})$ is the distribution function for the quasiparticles. The dynamics of the momentogenesis is determined by the spectral-flow.

Another property of the vortex is that it can change the topological charge of the vacuum. The vortex thus serves as the instanton, if the transition between the vacua of different topological charges occurs by tunneling, or as the sphaleron, if the transition occurs by thermal activation. In the Weinberg-Salam model the transition between topologically different vacua is accompanied by a change of the baryonic charge. The same happens with the fermionic charge of the superfluid. In the torus geometry the relevant fermionic charge of the vacuum is the momentum L_z, while the topological charge is the winding number of the phase Φ along the channel:

$$L_z(vac) = \hat{\mathbf{z}} \cdot \int d^3r \ \mathbf{r} \times (\rho \mathbf{v}_s) \ \ , \ \ N(vac) = \frac{1}{\kappa} \oint d\mathbf{r} \cdot \mathbf{v}_s \ \ . \tag{4.1.2}$$

These two charges are related:

$$\frac{L_z(vac)}{\hbar} = \frac{1}{2}\mathcal{N}(vac)N(vac) \ \ , \ \ \mathcal{N}(vac) = \frac{\rho}{m_3}V_{\rm torus} \ . \tag{4.1.3}$$

Here $\mathcal{N}(vac)$ is the total number of the ^{3}He atoms in the vacuum within the torus volume $V_{\rm torus}$. Note that $L_z(vac)/\hbar$ is always integral, even if the total number of ^{3}He atoms is odd. In the latter case one atom has no partner to form a Cooper pair and thus is not in the vacuum even at $T = 0$: it belongs to the *matter*. Another example of the nonzero fermionic charge of the *matter* even in the ground state will be discussed in Sec.5.1.

The change of the topological charge of the vacuum occurs by nucleation of a segment of the vortex with winding number N_v on the surface of the channel. The vortex sweeps the cross section of the channel and then disappears on the surface. In this process the topological charge $N(vac)$ changes by N_v. The superflow velocity in the channe, $v_s = \kappa N(vac)/L$, decreases by the value $\Delta v_s = \kappa N_v/L$ (where L is the length of the annular channel) which leads to the change of the fermionic charge of the vacuum by $\Delta L_z(vac)/\hbar = \frac{1}{2}\mathcal{N}(vac)N_v$. In electroweak theory this corresponds to the relation between the change of topological number of the vacuum (Chern-Simons number N_{CS}) and the change of the baryonic charge ΔB:

$$\Delta B = N_F \Delta N_{CS} \ , \tag{4.1.5}$$

where N_F is the number of the fermionic families. The difference from condensed matter case is that in the relativistic theories with local gauge symmetry the energy of the vacuum does not depend on its topological quantum number, while in the case of superfluid/superconductor the energy of the vacuum is $\propto v_s^2 \propto N^2(vac)$.

3.2. Quantum nucleation. Instanton.

The instanton is the quantum tunneling between two vacua, which is realized by the quantum nucleation of the vortex loop. At low T the spectral flow is suppressed and the tunneling is determined by the classical action which kinetic term is related to the Magnus force:

$$S = \kappa N_v \rho V + \int dt\ E \quad . \tag{4.2.1}$$

Here E is the energy of the vortex line. The first term is the kinetic part and is an analog of the Wess-Zumino action, where V is the volume bounded by the area swept by the vortex loop between nucleation and annihilation.[25] Variation of this term leads to the Magnus force. This term is of topological origin and leads to quantization of the total number number of particles in the vacuum, $\mathcal{N}(vac) = (\rho/m)V_{total}$, where V_{total} is the total volume of the vessel. The quantization occurs due to the ambiguity in choosing the volume swept by the loop: it can be V or $V - V_{total}$. Since the exponent $e^{iS/\hbar}$ should not depend on this choice, the difference between the actions, $\Delta S = m\kappa N_v \mathcal{N}(vac)$, must be multiple of $2\pi\hbar$. For ^{4}He, where $\kappa = 2\pi\hbar/m$ this gives an integral value of $\mathcal{N}(vac)$, while for ^{3}He, where $\kappa = \pi\hbar/m$, the number of particles in the vacuum must be even. The volume law for the action is the property of the global vortices where the field $\mathbf{v}_s$ generated by the vortex line is not screened by the gauge field.

In quantum tunnelling the action along the instanton trajectory contains an imaginary part, which gives the nucleation rate $\Gamma \propto e^{-2{\rm Im}S/\hbar}$. Let us consider the large $N(vac)$ case, when the velocity in the annular channel, $v_s = \kappa N(vac)/L$, is large and the nucleation occurs close to the pinning center at the wall of container. For the hemispherical pinning center the instanton is the trajectory of the vortex half-ring coaxial with the pinning center and oriented perpendicular to the flow (axis x). Far from the pinning center the energy of the vortex loop of radius r is

$$E = E_0(r) + \mathbf{v}_s \cdot \mathbf{p}(r)\ ,\ E_0(r) = \frac{1}{2}\rho N_v^2 \kappa^2 r \ln\frac{r}{\xi}\ ,\ p_x(r) = \pi\rho N_v \kappa r^2 \tag{4.2.2}$$

where $E_0(r)$ is the line tension, ξ is the core radius, and $\mathbf{p}(r) = \hat{\mathbf{x}} p_x$ is the linear momentum of the loop. The vortex nucleated from the vacuum state must have zero energy, $E = 0$. This determines the radius of the nucleated vortex:

$$R_0 = \frac{\kappa N_v}{2\pi v_s} \ln\frac{R_0}{\xi} \quad . \tag{4.2.3}$$

The calculation of the instanton action for the ^{4}He vortex with $N_v=1$ shows that it is proportional to the number $\mathcal{N}_0$ of particles involved in the process of tunneling:[26]

$$\mathrm{Im}S = 2\pi\hbar\mathcal{N}_0 \ \ , \ \ \mathcal{N}_0=\frac{\rho}{m}\frac{2\pi}{3}R_0^3 \ . \tag{4.2.4}$$

For the ^{3}He case and for the vortex tunneling in superconductors, the situation can be different because of the fermion zero modes in the vortex core, which lead to the spectral flow. This question is still open.

3.3. Thermal activation. Sphaleron

The thermal activation, which can occur at nonzero T, is determined by the so called sphaleron[7] – the saddle-point stationary solution, which is intermediate between vacuum states with different topological number $N(vac)$. In superfluids the sphaleron is represented by a metastable vortex loop, stationary in the heat-bath frame: its velocity $dE/d\mathbf{p}=0$, which corresponds to the maximum of the Doppler shifted energy $E=E_0(r)+\mathbf{v}_s\cdot\mathbf{p}(r)$. Using the Eq.(4.2.2) for E one obtains the radius and the energy of the sphaleron

$$r_\circ=\frac{\kappa N_v}{4\pi v_s}\ln\frac{r_\circ}{\xi} \ , \ E_0(r_\circ)=\frac{1}{2}\rho\kappa^2N_v^2 r_\circ \ln\left(r_\circ/\xi\right) \ . \tag{4.3.1}$$

The sphaleron is created by thermal activation with a rate $\exp-(E_0(r_\circ)/T)$. After that the vortex ring grows spontaneously, thus decreasing its Doppler shifted energy E, until it disappears at the wall of the torus leaving behind the flow pattern with the topological number $N(vac)$ reduced by N_v.

4. DISCUSSION. OTHER SCENARIA.

The quantum field theory in condensed matter and in superfluid ^{3}He in particular can provide analogies for many different scenaria of baryogenesis. For example, the production of the linear momentum[27] and spin momentum[28] during the motion of the interface between the ^{3}He-A and ^{3}He-B can simulate the mechanisms related to the baryogenesis by domain walls. Here we briefly mention the analogy of baryogenesis by the baryonic charge condensate.[4]

4.1. Nonzero charge of *matter* induced by vacuum charge.

The vacuum with nonzero fermionic charge takes place in ferromagnets, where the charge is played by the discrete projection S_z of spin to the direction of the spontaneous magnetization. This charge is conserved above and below the broken symmetry phase transition. Since we are interested in the quantum field theory, let us consider the hypothetical phases of the superfluid ^{3}He, which have spontaneous vacuum spin $S_z(vac)$. This can be for example A$_1$-phase or β phase ((Eq.(6.53) in[29]), which could appear if the parameters of the systems were more favourable. The magnitude of the vacuum spin in such phases is

$$S_z(vac) \sim \hbar \mathcal{N}(vac) \frac{1}{(k_F \xi)^2} \quad , \tag{5.1.1}$$

where $\mathcal{N}(vac)$ is the total number of the ^{3}He atoms; $\xi = p_F/m\Delta$ is the superfluid coherence length, and Δ is the gap. If there are no boundaries and thus no exchange with environment, then $S_z(vac)$ should be compensated by the opposite amount of the spin of the quasiparticles. In this case the ground state is not the vacuum: it contains quasiparticles, with number

$$N_{qp} = 2\frac{S_z(vac)}{\hbar} \sim \mathcal{N}(vac) \frac{1}{(k_F \xi)^2} \quad . \tag{5.1.2}$$

Since $k_F\xi \sim 10^2 - 10^3$, the number of the particles in the *matter*, N_{qp}, is much smaller than the number of the particles in the vacuum, $\mathcal{N}(vac)$, and thus the vacuum is not disturbed by the *matter*. The energy of the *matter* is about ΔN_{qp} (or even less if the quasiparticles are massless), and this is essentially smaller than the vacuum energy.

The conservation of the total spin after the transition can be also provided by the formation of two domains with opposite direction of spin. This also costs energy: now it is the energy of the wall between two domains $E_{wall} \sim m_3 p_F \Delta^2 \xi V_{total}^{2/3}$. Comparing this to the energy of the quasiparticles, one finds that quasiparticles in the ground states are preferred over a domain wall if the size of the sample is small enough. The condition for the total number of ^{3}He atoms (the total number of the particles in the vacuum) is:

$$\mathcal{N}(vac) < (\xi k_F)^6 \quad . \tag{5.1.3}$$

The symmetry breaking in the Universe can also give rise to nonzero vacuum fermionic charge, the baryonic number $B(vac)$.[4] Then the matter acquires a baryonic charge opposite to the vacuum charge, $B_{matter} = -B(vac)$, if (1) the total baryonic charge is conserved; (2) if the baryonic exchange between the vacuum and matter is not suppressed; and (3) if the energy of the domain wall between matter and antimatter is larger than the energy of the excess baryonic matter. Many details of this scenario of baryogenesis can be modelled in condensed matter. The transition to superfluid ^{3}He-A in a small droplet of ^{3}He can lead to quasiparticles in the ground state, because the A-phase state has a nonzero vacuum value of the orbital momentum. The decay of the vacuum spin into the spin of the fermions is one of the heavily discussed topics in the so-called magnetic superfluidity of ^{3}He-B, where the coherent precession of the magnetization takes place with all the signatures of the spin condensate;[30] and so on.

The related problem is the formation of the galactic magnetic fields. In a recent paper Joyce and Shaposhnikov[31] suggested a new scenario: the galactic magnetic fields are generated from the primordial hypercharge magnetic fields. The latter are spontaneously formed by the right-handed electrons through the Abelian anomaly. It appears that the process of generation of hypercharge magnetic field from the fermionic charge of matter is described by the same equations as the process of the superflow instability in ^{3}He-A, which was intensively discussed theoretically and has been recently investigated experimentally.[32] Both processes are governed by the chiral anomaly. In ^{3}He-A, the motion of the normal component, which contains the fermionic charge – the linear momentum of quasiparticles, is unstable towards the inhomogeneous condensate – the texture of the $\hat{\mathbf{l}}$-vector, which corresponds to the hypercharge magnetic field.

In conclusion, when the problem of the baryogenesis is finally solved, it will appear that the key to the problem has been provided by condensed matter physics.

REFERENCES

1. G.Volovik, T.Vachaspati, *Int. J. Mod. Phys.* **B 10**, 471 (1996).
2. S. Adler, *Phys. Rev.* **177**, 2496 (1969).
3. J.S. Bell, R. Jackiw, *Nuovo Cim.* **A 60**, 47 (1969).
4. A.D. Dolgov, *Phys. Rep.* **222**, 310 (1992).
5. A.Vilenkin, E.P.S.Shellard, *Cosmic Strings and Other Topological Defects*, (Cambridge University Press,1994).
6. M.B. Hindmarsh and T.W.B. Kibble, *Phys. Rep.* **58**, 477 (1995).
7. N. Turok, in *Formation and Interactions of the Topoogical Defects*, eds. A.C. Davis and R. Brandenberger, Plenum Press, New York and London, 1995.
8. T. Vachaspati, G.B. Field, *Phys. Rev. Lett.* **73**, 373 (1994); **74**, *Errata* (1995).
9. J. Garriga and T. Vachaspati, *Nucl. Phys.* **B438**, 161 (1995).
10. G.Volovik, *Exotic Properties of Superfluid ^{3}He*, (World Scientific,Singapore,1992).
11. G.E. Volovik, *JETP* **75**, 990 (1992).
12. N.B. Kopnin, *Phys. Rev.* **B 47**, 14354 (1993).
13. E.B. Sonin, *Phys. Rev.* **B 55**, 485 (1997).
14. T.D.C. Bevan, A.J. Manninen, J.B. Cook, *et al*, *Nature* **386**, 689 (1997)
15. T.D.C. Bevan, A.J. Manninen, J.B. Cook, *et al*, to be published.
16. C.Caroli, *et al*, *Phys. Lett.* **9** (1964) 307.
17. G.E. Volovik, *ZhETF* **104**, 3070 (1993) [*JETP* **77**, 435 (1993)].
18. M. Stone, *Phys. Rev.* **B 54**, 13222 (1996); cond-mat/9708017.

19. N.B. Kopnin, G.E. Volovik, and Ü. Parts, *Europhys. Lett.* **32***, 651 (1995).*
20. G.E. Volovik and V.P. Mineev, *JETP* **56**, 579 (1982).
21. N.B. Kopnin, V.E. Kravtsov, *ZhETF*, **71**, 1644 (1976); [*JETP*, **44**, 861 (1976)].
22. N.B. Kopnin and G.E. Volovik, *Phys. Rev. Lett.* **79**, 1377 (1997).
23. N.B. Kopnin and G.E. Volovik, cond-mat/9706082.
24. T. Jacobson and G.E. Volovik, to be published.
25. M. Rasetti and T. Regge, *Physica* **A 80**, 217 (1975).
26. G.E. Volovik, *Pis'ma ZhETF* **15**, 116 (1972) [*JETP Lett.* **15**, 81 (1972)].
27. S. Yip, *Phys. Rev.* **B 35**, 8733 (1987).
28. N. Schopohl and D. Waxman, *Phys. Rev. Lett.* **63**, 1696 (1989).
29. D. Vollhardt, P. and P. Wölfle *The superfluid phases of helium 3*, Taylor and Francis, London - New York - Philadelphia, 1990.
30. Yu.M. Bunkov, in *Progress in Low Temperature Physics*, ed. W. Halperin, Elsevier, **14**, 69 (1995).
31. M. Joyce and M. Shaposhnikov, *Phys. Rev. Lett.*,**79**, 1193 (1997).
32. V.M.H. Ruutu, J. Kopu, M. Krusius, U. Parts, B. Plaais, E.V. Thuneberg, and W. Xu, "Critical Velocity in Rotating Superfluid 3He-A", preprint LTL-E97024 (http://boojum.hut.fi/abstracts/lowtemp/LTL-E97024.html), submitted to *Phys. Rev. Lett.*.

Figure Captions:

Fig.1 Connection of ^{3}He with the rest of physics.

Fig.2 Right electron in magnetic field. The lowest ($n = 0$) Landau level is asymmetric. The motion of the particles along the anomalous branch leads to the creation of particles from the Dirac vacuum.

Fig.3 Asymmetric branch in the spectrum of the fermions bound to the core of the vortex in s-wave superconductor. The motion of the particles along the anomalous branch leads to the spectral-flow force acting on the vortex.



Alice string
Witten string
Torsion string
Z-string
Monopoles
on string
String on wall

Symmetry breaking (anisotropy of vacuum)
Higgs field
Fermionic and bosonic excitations
Screening - antiscreening
Vacuum instability in strong field
Fermionic charge of vacuum
Axial anomaly

Baryogenesis
Kibble
mechanism

cosmic strings

physical vacuum

Vortex dynamics
Gap nodes
Low -T scaling
1/2-vortex

Cosmological
constant
Effective gravity
Bi-metric gravity

cosmology

High Energy Physics

high-T super-conductivity

QHE
Quantum
statistics
of skyrmions

vacuum gravity

Gravity

^{3}He

Condensed Matter

2D systems

black holes

Nuclear Matter

Plasma Physics

super-fluidity

Horizon
Hawking radiation
Information puzzle

Bose condensate
Magnetic
superfluidity

Coulomb
plasma

neutron stars

nuclear physics

hydro-dynamics

Superfluidity
of neutron star
Vorticies
Glitches

Nuclei vs ^{3}He droplet
Shell model
Pair-correlations
Collective modes

General
Multi-fluid
Relativistic
Hydrodynamics
of rotating superfluid
Magnetohydrodynamic



$E(p_z)$
n=2
n=1
n=0
e_R
p_z
n=-1
n=-2


E (L$_z$, p$_z$ = 0)
1/2
3/2
5/2
7/2
L$_z$